

Approximate McKean-Vlasov Representations for a class of SPDEs*

D. Crisan[†] J. Xiong[‡]

22 December 2004

Abstract

The solution $\vartheta = (\vartheta_t)_{t \geq 0}$ of a class of linear stochastic partial differential equations is approximated using Clark's robust representation approach ([1], [2]). The ensuing approximations are shown to coincide with the time marginals of solutions of a certain McKean-Vlasov type equation. We prove existence and uniqueness of the solution of the McKean-Vlasov equation. The result leads to a representation of ϑ as a limit of empirical distributions of systems of equally weighted particles. In particular, the solution of the Zakai equation and that of the Kushner-Stratonovitch equation (the two main equations of nonlinear filtering) are shown to be approximated the empirical distribution of systems of particles that have equal weights (unlike those presented in [15] and [16]) and do not require additional correction procedures (such as those introduced in [5], [6], [10], etc).

Keywords: Stochastic Partial Differential Equations, particle approximations, McKean-Vlasov equations, Zakai equation, Kushner-Stratonovitch equation, nonlinear diffusions.

MSC 2000: 60H15, 60K35, 35R60, 93E11.

*First draft 1 October 2003, current version 22 December 2004.

[†]Department of Mathematics, Imperial College London, 180 Queen's Gate, London SW7 2BZ, UK.

[‡]Department of Mathematics, University of Tennessee, Knoxville, TN 37996-1300 USA. Research supported partially by NSA.

1 Introduction

Let $(\Omega, \mathcal{F}, P)$ be a probability space on which we have defined an m -dimensional Wiener process $W = (W^i)_{i=1}^m$. Let $M(\mathbb{R}^d)$ be the space of finite Borel measures defined on the d -dimensional Euclidean space $\mathbb{R}^d$ and $(\vartheta_t)_{t \geq 0}$ be an $M(\mathbb{R}^d)$ -valued stochastic process satisfying the following linear stochastic partial differential equation

$$\vartheta_t(\varphi) = \vartheta_0(\varphi) + \int_0^t \vartheta_s (\alpha_s \varphi + L_s \varphi) ds + \sum_{j=1}^m \int_0^t \vartheta_s (\beta_s^j \varphi) dW_s^j, \quad (1)$$

where $L = (L_s)_{s \geq 0}$, $L_s : C_b(\mathbb{R}^d) \rightarrow C_b(\mathbb{R}^d)$ is the second order elliptic differential operator

$$L_s \varphi = \frac{1}{2} \sum_{i,j=1}^d a_s^{ij} \partial_{x_i} \partial_{x_j} \varphi + \sum_{i=1}^d b_s^i \partial_{x_i} \varphi, \quad (2)$$

and φ is a function in the domain of L , $\varphi \in \bigcap_{s \geq 0} \mathcal{D}(L_s)$. For simplicity, we will assume that ϑ_0 is a probability measure. If $\alpha \equiv 0$, then (1) is called the Zakai or the Duncan-Mortensen-Zakai equation (cf [11], [21], [27]). The Zakai equation has been studied extensively over the last 40 years because of its importance in non-linear filtering theory (see [20], [22]): In non-linear filtering, $\bar{\vartheta}_t$, the normalized form of ϑ_t , is the conditional distribution of a (non-homogeneous) Markov process ξ_t with infinitesimal generator L given the observation process W which satisfies the evolution equation

$$W_t = \int_0^t \beta_s(\xi_t) ds + V_t, \quad (3)$$

In (3), V is an m -dimensional Wiener process independent of ξ . Within the filtering problem W is not a Brownian motion as it is assumed in the current set-up. However, it becomes a Brownian motion after a suitable change of measure (Girsanov transformation). One can prove that $\bar{\vartheta}_t$ satisfies the following (non-linear) stochastic partial differential equation

$$\begin{aligned} \bar{\vartheta}_t(\varphi) &= \bar{\vartheta}_0(\varphi) + \int_0^t \bar{\vartheta}_s (\alpha_s \varphi + L_s \varphi) - \bar{\vartheta}_s(\alpha_s) \bar{\vartheta}_s(\varphi) ds \\ &\quad + \sum_{j=1}^m \int_0^t \bar{\vartheta}_s (\beta_s^j \varphi) - \bar{\vartheta}_s(\beta_s^j) \bar{\vartheta}_s(\varphi) (dW_s^j - \bar{\vartheta}_s(\beta_s^j) ds), \end{aligned} \quad (4)$$

If $\alpha \equiv 0$, then equation (4) is called the Fujisaki-Kallianpur-Kunita or Kushner-Stratonovitch equation (cf. [12], [17]). In general, neither (1) nor (4) have explicit solutions, though one can approximate them by numerical means. As expected, there is a wide variety of methods to do this (see, for example, [3] and the references therein). Among them, particle methods seem to be in many cases quite effective. The starting point for a particle approximation of ϑ_t is the following Feynman-Kač representation

$$\vartheta_t(\varphi) = E[\varphi(X_t) A_t(X) | W], \quad (5)$$

where $X = (X_t)_{t \geq 0}$ is a Markov process with infinitesimal generator L independent of W with initial distribution $\bar{\vartheta}_0$ and $A_t(X)$ is defined as

$$A_t(X) = \exp \left(\int_0^t \alpha_s(X_s) ds + \sum_{j=1}^m \left(\int_0^t \beta_s^j(X_s) dW_s^j - \frac{1}{2} \int_0^t \beta_s^j(X_s)^2 ds \right) \right), \quad t \geq 0.$$

Since in (5) X is independent of W , one can use the Monte Carlo method to compute ϑ_t . That is, ϑ_t has the representation

$$\vartheta_t = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n A_t(X^i) \delta_{X_t^i}, \quad (6)$$

where $X^i, i > 0$ are independent realizations of X . In other words, $\{X_t^i, A_t(X^i)\}_{i=1}^\infty$ satisfies the following system of SDEs

$$\begin{cases} dX_t^i &= b_t(X_t^i) dt + \sigma_t(X_t^i) dB_t^i \\ dA_t(X^i) &= A_t(X^i) \alpha_t(X_t^i) dt + \sum_{j=1}^m A_t(X^i) \beta_t^j(X_t^i) dW_t^j \end{cases} \quad (7)$$

where $B^i, i > 0$, are mutually independent d -dimensional Brownian motions independent of W and $\sigma_s = (\sigma_s)_{i,j=1}^d$ is chosen to satisfy $a_s = \sigma_s \sigma_s^\top$ ($\sigma_s^\top$ is the transpose of σ_s). As demonstrated in [15] and [16], representations of the type (5) and (6) hold true for a far wider class of SPDEs than the one described by (1).

However, the convergence in (6) is very slow. That is because the variance of the weights $A_t(X^i), i > 0$, increases exponentially fast with time. The effect is that most of weights decrease to 0 with only a few becoming very large. In order to offset this outcome, a wealth of methods have been proposed. In filtering theory, the generic name for such a method is that of a *particle filter* ([4], [5], [6], [7], [8], [10], etc.). The standard remedy is to introduce an additional procedure that removes particles with small weights and adds additional particles in places where the existing one have large weights. Put differently, one applies at certain times a

branching procedure by which, each particle will be replaced by a random number of “offsprings” with a mean proportional with its corresponding weight. This branching procedure is a double edge sword: applying it too often may actually decrease the rate of convergence (cf. [9]).

In this paper, we seek a different remedy to the slow convergence of the Monte Carlo method. Heuristically we seek to keep the weights of the particles equal without introducing an additional procedure but only by amending the motion of the particles in a way that will take into account the state of the entire system. This will be achieved in two steps:

First we will show that ϑ_t and its normalised version $\bar{\vartheta}_t$ admit a robust version following the approach first introduced by Clark ([1], see also [2]). By this we mean that ϑ_t and, respectively, $\bar{\vartheta}_t$ depends continuously on the generating Brownian path $s \rightarrow W_s(\omega)$, $\omega \in \Omega$. Hence if we consider an approximating sequence of paths $s \rightarrow W_s^\delta(\omega)$ that will converge to $s \rightarrow W_s(\omega)$ as δ tends to 0, then the corresponding measure valued processes ϑ_t^δ and $\bar{\vartheta}_t^\delta$ will converge to ϑ_t and, respectively, $\bar{\vartheta}_t$ with a certain rate of convergence $r(t, W(\omega), \delta)$.

The second step is to prove that $\bar{\vartheta}_t^\delta$ has the asymptotic representation

$$\bar{\vartheta}_t^\delta = \lim_{n \rightarrow \infty} \bar{\vartheta}_t^{\delta, n}, \quad \text{where } \bar{\vartheta}_t^{\delta, n} = \frac{1}{n} \sum_{i=1}^n \delta_{X_t^{i, \delta}}, \quad (8)$$

and X^i , $i \geq 0$ are *non-linear diffusions* satisfying the non-linear SDE

$$dX_t^{i, \delta} = \tilde{b}_t^\delta(\tilde{\vartheta}_t^\delta, X_t^{i, \delta}) dt + \sigma_t(X_t^{i, \delta}) dB_t^i. \quad (9)$$

In (9), $\tilde{\vartheta}_t^\delta$ is the conditional distribution of $X_t^{i, \delta}$, B^i are mutually independent Brownian motions and the function $\tilde{b}_t^\delta$ will depend intrinsically on the chosen path $s \rightarrow W_s(\omega)$. We will show that $\tilde{\vartheta}_t^{\delta, n} = \bar{\vartheta}_t^{\delta, n}$ and that, by suitably choosing the parameter $\delta = \delta(W(\omega), n)$, we will have

$$\bar{\vartheta}_t = \lim_{n \rightarrow \infty} \bar{\vartheta}_t^{\delta, n}, \quad \text{where } \bar{\vartheta}_t^{\delta, n} = \frac{1}{n} \sum_{i=1}^n \delta_{X_t^{i, \delta(W(\omega), n)}}.$$

Hence $\bar{\vartheta}_t$ has an asymptotic representation involving particles with equal weights. Finally in order to obtain the corresponding asymptotic representation for ϑ_t we need to “unnormalize” $\bar{\vartheta}_t^n$, i.e., to attach to each particle a weight a_t^n (the same one) which converges to $\vartheta_t(1)$ as n tends to infinity. For example, one could choose

$$a_t^n = \exp \left(\int_0^t \bar{\vartheta}_s^{\delta, n}(\alpha_s) ds + \sum_{j=1}^m \left(\int_0^t \bar{\vartheta}_s^{\delta, n}(\beta_s^j) d(W_s^\delta)^j - \frac{1}{2} \int_0^t \bar{\vartheta}_s^{\delta, n}(\beta_s^j)^2 ds \right) \right).$$

Hence, we obtain the asymptotic representation ϑ_t which uses systems of particles with equal weights.

$$\vartheta_t = \lim_{n \rightarrow \infty} \frac{a_t^n}{n} \sum_{i=1}^n \delta_{X_t^{i, \delta(W \cdot (\omega), n)}}.$$

We remark that we are not aware of a similar result even for the simpler case when (1) has no stochastic term, i.e., (1) is a second order elliptic PDE with a zero-order term. In this case the first step described above is not required: One obtains directly an asymptotic representation of the solution

$$\bar{\vartheta}_t = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta_{X_t^i}$$

where the X^i , $i \geq 0$ are non-linear diffusions satisfying a non-linear SDE of the type (9).

2 The robust representation of $\bar{\vartheta}_t$

In this section we introduce the robust representation formula for ϑ_t and, respectively, $\bar{\vartheta}_t$. By robustness here we mean that the dependence of the generating Brownian path $t \rightarrow W_t(\omega)$ is continuous. The formula is similar to, and inspired by, the robust version of the integral representation formula of nonlinear filtering as presented in [1] and [2]. At a formal level, the formula is derived by a process of integration-by-parts applied to the stochastic integrals that appear in the Feynman-Kač representation (5). The rigorous justification of the formula is identical with that of the corresponding result in [2] and for this reason we omit it here.

For the existence of the robust representation of ϑ_t and, respectively, of $\bar{\vartheta}_t$ we follow Theorems 1 and 2 in [2]. For this we need to impose the following conditions:

RR1: For all $j = 1, \dots, d$ we assume that $(s, x) \rightarrow \beta_s^j(x) \in C_{1,2}(\mathbb{R}_+ \times \mathbb{R}^d)$, hence $s \rightarrow \beta_s^j(X_s)$ is a semimartingale (we denote by $\beta^j(X)^m$ its martingale part and by $\beta^j(X)^{fv}$ its finite variation part) such that, for all $k > 0$ and all $t > 0$

$$E \left[\exp \left(k \sum_{j=1}^d \int_0^t d \langle \beta^j(X)^m \rangle_s \right) \right] < \infty, \quad E \left[\exp \left(k \sum_{j=1}^d \int_0^t |d \beta^j(X)^{fv}| \right) \right] < \infty.$$

RR2: For all $t > 0$ we have

$$E \left[\exp 2 \int_0^t \alpha_s(X_s) ds \right] < \infty.$$

Let $y. = \{y_s, s \geq 0\} \in C_{\mathbb{R}^d}([0, \infty))$ be a continuous path $\mathbb{R}^d$ and define $\Theta^{y.} = (\Theta_t^{y.})_{t \geq 0}$ to be the following $M(\mathbb{R}^d)$ -valued stochastic process (φ is a bounded Borel measurable function)

$$\begin{aligned} \Theta_t^{y.}(\varphi) = E \left[\varphi(X_t) \exp \left(\int_0^t \alpha_s(X_s) ds + \sum_{j=1}^m \beta_s^j(X_s) y_t^j \right. \right. \\ \left. \left. - \sum_{j=1}^m \left(I^j(y.) + \frac{1}{2} \int_0^t \beta_s^j(X_s)^2 ds \right) \right) \right], \end{aligned}$$

where $X = (X_t)_{t \geq 0}$ is a Markov process with infinitesimal generator L independent of W with initial distribution $\bar{\nu}_0$ and $I^j(y.)$ is a version of the stochastic integral $\int_0^t y_s^j d\beta_s^j(X_s)$. Let $\bar{\Theta}^{y.} = (\bar{\Theta}_t^{y.})_{t \geq 0}$ be normalized version of the $M(\mathbb{R}^d)$ -valued stochastic $\bar{\Theta}^{y.} = (\bar{\Theta}_t^{y.})_{t \geq 0}$

$$\bar{\Theta}_t^{y.} = \frac{\Theta_t^{y.}}{\Theta_t^{y.}(\mathbf{1})},$$

where $\mathbf{1}$ is the constant function $\mathbf{1}(x) = 1$ for all $x \in \mathbb{R}^d$ ($\Theta^{y.}(\mathbf{1})$ is the mass process associated to $\Theta^{y.}$). Following Theorem 1 in [2], under the conditions **RR1** and **RR2**, $\Theta^{y.}$ and $\bar{\Theta}^{y.}$ depend continuously on the path $y.$. Moreover, for any $R > 0$, there exist two constants $K = K(R, t)$ and $\bar{K} = \bar{K}(R, t)$ such that

$$\begin{aligned} |\Theta_t^{y.^1}(\varphi) - \Theta_t^{y.^2}(\varphi)| &\leq K \|\varphi\| \|y.^1 - y.^2\|_t \\ |\bar{\Theta}_t^{y.^1}(\varphi) - \bar{\Theta}_t^{y.^2}(\varphi)| &\leq \bar{K} \|\varphi\| \|y.^1 - y.^2\|_t \end{aligned} \quad (10)$$

for any bounded measurable function φ ($\|\varphi\| \triangleq \sup_{x \in \mathbb{R}^d} |\varphi(x)| < \infty$) and for any two paths $y.^1, y.^2$ such that $\|y.^1\|_t, \|y.^2\|_t \leq R$ (where $\|\cdot\|_t$ is defined as $\|\alpha\|_t \triangleq \max_{1 \leq i \leq d} \max_{s \in [0, t]} |\alpha_s^i|$). Furthermore, if we use the norm $\|\cdot\|_w$ on the set of finite signed measures $M^s(\mathbb{R}^d)$

$$\|\mu\|_w = \sup_{\{\varphi \in C_b(\mathbb{R}^d) \mid \|\varphi\| \leq 1\}} |\mu(\varphi)|,$$

then, from (10), we deduce that

$$\left\| \Theta_t^{y.^1} - \Theta_t^{y.^2} \right\|_w \leq K \|y.^1 - y.^2\|_t, \quad \left\| \bar{\Theta}_t^{y.^1} - \bar{\Theta}_t^{y.^2} \right\|_w \leq \bar{K} \|y.^1 - y.^2\|_t. \quad (11)$$

Following Theorem 2 in [2], $\Theta^{W.}$ and $\bar{\Theta}^{W.}$ are the robust versions of ϑ and, respectively, $\bar{\vartheta}$. More precisely for all $t \geq 0$,

$$\Theta_t^{W.} = \vartheta_t \text{ and } \bar{\Theta}_t^{W.} = \bar{\vartheta}_t, \text{ } P\text{-almost surely} \quad (12)$$

and the null set can be taken to be independent of t since all processes involved are time-continuous.

The robust representation result also enables us to fix the generating Brownian path $s \rightarrow W_s(\omega)$ and proceed to approximate the corresponding $\bar{\vartheta}_t^\delta$ and, implicitly, $\bar{\vartheta}_t$ for this fixed, but arbitrary, path $s \rightarrow W_s(\omega)$.

We will replace the generating Brownian path $t \rightarrow W_t(\omega)$ with a smoother one. For example, we choose a partition of the time interval $[0, \infty)$ of the form

$$0 = t_0 < t_1 < \dots < t_i < \dots$$

where $t_i = i\delta$, $i \geq 0$ and define the following piecewise linear path $t \rightarrow W_t^\delta(\omega)$

$$W_t^\delta(\omega) = W_{t_i}(\omega) + \frac{W_{t_{i+1}}(\omega) - W_{t_i}(\omega)}{\delta} (t - t_i), \quad \text{for } t \in [t_i, t_{i+1}).$$

Let $\vartheta_t^\delta \triangleq \Theta_t^{W^\delta}$ and $\bar{\vartheta}_t^\delta \triangleq \bar{\Theta}_t^{W^\delta}$. Then, from (11), it follows that,

$$\lim_{\delta \rightarrow 0} \vartheta_t^\delta = \vartheta_t \quad \text{and} \quad \lim_{\delta \rightarrow 0} \bar{\vartheta}_t^\delta = \bar{\vartheta}_t,$$

where the above convergence is in the weak topology on $M(\mathbb{R}^d)$. Moreover, we have the following rates of convergence: there exist two constants $K = K(\varpi_t(\omega), t)$ and $\bar{K} = \bar{K}(\bar{\varpi}_t(\omega), t)$ where $\varpi_t(\omega) = \sup_{s \in [0, t+\delta]} \max_{1 \leq i \leq d} |W_s^i(\omega)|$ such that

$$\|\vartheta_t^\delta - \vartheta_t\|_w \leq K \varpi_t^\delta(\omega), \quad \|\bar{\vartheta}_t^\delta - \bar{\vartheta}_t\|_w \leq \bar{K} \bar{\varpi}_t^\delta(\omega)$$

and $\varpi_t^\delta(\omega) = \max_{j=0, [\delta]} \sup_{t \in [t_i, t_{i+1}]} \max_{1 \leq i \leq d} |W_t^i(\omega) - W_{t_j}^i(\omega)|$. Moreover since $s \rightarrow W_s(\omega)$ is continuous, we can choose $\delta = \delta(\omega, n)$, so that $\varpi_t^\delta(\omega) \leq \frac{1}{n}$. Hence

$$\begin{aligned} \left\| \vartheta_t^{\delta(\omega, n)} - \vartheta_t \right\|_{\mathcal{M}} &\leq \left\| \vartheta_t^{\delta(\omega, n)} - \vartheta_t \right\|_w \leq \frac{K}{\sqrt{n}}. \\ \left\| \bar{\vartheta}_t^{\delta(\omega, n)} - \bar{\vartheta}_t \right\|_{\mathcal{M}} &\leq \left\| \bar{\vartheta}_t^{\delta(\omega, n)} - \bar{\vartheta}_t \right\|_w \leq \frac{\bar{K}}{\sqrt{n}}. \end{aligned} \tag{13}$$

In (13), the norm $\|\cdot\|_{\mathcal{M}}$ ¹ is defined as

$$\|\mu\|_{\mathcal{M}} \triangleq \sum_{k=0}^{\infty} \frac{|\mu(\varphi_k)|}{2^k \|\varphi_k\|}, \quad \mu \in M^s(\mathbb{R}^d), \tag{14}$$

where $\varphi_0 = 1$ and $(\varphi_k)_{k>0}$ are the elements of a dense set $\mathcal{M} \in C_b(\mathbb{R}^d)$. This norm will be used to prove convergence in the next step of the construction.

¹Both norms $\|\cdot\|_w$ and $\|\cdot\|_{\mathcal{M}}$ induce the weak topology on $M(\mathbb{R}^d)$.

It is easy to see that ϑ_t^δ satisfies the following linear PDE (written in weak form)

$$\vartheta_t^\delta(\varphi) = \vartheta_0(\varphi) + \int_0^t \vartheta_s^\delta (\alpha_s^\delta \varphi + L_s \varphi) ds \quad (15)$$

where

$$\alpha_s^\delta = \alpha_s - \frac{1}{2} \sum_{j=1}^m \left((\beta_s^j)^2 + \beta_s^j \frac{W_{t_{i+1}}^i(\omega) - W_{t_i}^i(\omega)}{\delta} \right), \quad s \in [t_i, t_{i+1}) \quad (16)$$

whilst $\bar{\vartheta}_t^\delta$ satisfies the following nonlinear PDE (again, written in weak form)

$$\bar{\vartheta}_t^\delta(\varphi) = \bar{\vartheta}_0(\varphi) + \int_0^t \bar{\vartheta}_s^\delta (\alpha_s^\delta \varphi + L_s \varphi) - \bar{\vartheta}_s^\delta (\alpha_s^\delta) \bar{\vartheta}_s^\delta (\varphi) ds \quad (17)$$

Note that, from (15), $\vartheta_t^\delta(1) = \exp \left(\int_0^t \bar{\vartheta}_s^\delta (\alpha_s^\delta) ds \right)$ (since $\vartheta_0^\delta(1) = 1$).

We introduce now additional conditions on the coefficients of the SPDE (1) to insure it has a unique solution and that the solution has a density with respect to the Lebesgue measure and its density is strictly positive everywhere.

EU: We will assume that the functions $\alpha_s, \beta_s^j, j = 1, \dots, m$, are bounded (with a bound independent of s). $a_s^{ij}, i, j = 1, \dots, d, b_s^i, i = 1, \dots, d$ are bounded (with a bound independent of s) and Lipschitz (with a Lipschitz constant independent of s). We also assume that the operator L is uniformly elliptic. That is there is a constant b such that for all times $s \geq 0$ and vectors $\xi = (\xi_i)_{i=1}^d \in \mathbb{R}^d$, we have

$$\langle \xi, a_s \xi \rangle \geq b \langle \xi, \xi \rangle,$$

where $\langle \cdot, \cdot \rangle$ is the standard inner product on $\mathbb{R}^d$. Further, we will assume that ϑ_0 has finite second moment.

PD: We will assume that ϑ_0 is absolutely continuous with respect to the Lebesgue measure and its density is strictly positive everywhere.

Assuming condition **EU** ensures that the SPDE (1) (see for example Jie&Kurtz) and the PDEs (15) and (17) have unique solutions. Further, the system of equations (7) and, in particular, the SDE satisfied by the processes X_i (which appear in the asymptotic representation (6) of ϑ_t) has a unique solution. In particular this ensures the existence of a Markov process $X = (X_t)_{t \geq 0}$ with infinitesimal generator L and initial distribution $\bar{\vartheta}_0$. Assuming **EU+PD** guarantees that, for any $t \geq 0$ the distribution of X_t is absolutely continuous with respect to the Lebesgue measure and its density is strictly positive everywhere.

Let us note that, from the Feynman-Kač representation (5), ϑ_t (and hence its normalized version $\bar{\vartheta}_t$) has the same support as the distribution of X_t . Similarly, ϑ_t^δ has the following representation

$$\vartheta_t^\delta(\varphi) = E \left[\varphi(X_t) \exp \left(\int_0^t \alpha_s^\delta(X_s) ds \right) \right]. \quad (18)$$

So, both ϑ_t^δ and $\bar{\vartheta}_t^\delta$ have the same support as the distribution of X_t , too. It follows from the above that ϑ_t , $\bar{\vartheta}_t$, ϑ_t^δ and $\bar{\vartheta}_t^\delta$ are all absolutely continuous with respect to the Lebesgue measure and their density is strictly positive everywhere. In the following we will denote by $x \rightarrow \bar{\vartheta}_t(x)$ and $x \rightarrow \bar{\vartheta}_t^\delta(x)$ the density of $\bar{\vartheta}_t$ and respectively $\bar{\vartheta}_t^\delta$ with respect to the Lebesgue measure.

3 The non-linear process

We introduce next the non-linear SDEs satisfied by the non-linear diffusions $X^{i,\delta}$, $i \geq 0$ appearing in the asymptotic representation (8) of $\bar{\vartheta}_t^\delta$. For this we need to define first the drift coefficient of the non-linear SDE (9). In the following, we fix the generating Brownian path $t \rightarrow W_t(\omega)$. The coefficient α_s^δ obviously depends on the path, however we will not make the dependence explicit. For this fixed path, the non-linear diffusions $X^{i,\delta}$ will be defined on a new probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$.

For arbitrary $f \in \mathcal{B}(\mathbb{R}^d)$ and $\mu \in \mathcal{P}(\mathbb{R}^d)$, define $f^\mu \in \mathcal{B}(\mathbb{R}^d)$ to be the function $f^\mu \triangleq f - \mu(f)$ and let $\Lambda_\mu f = (\Lambda_\mu^j f)_{j=1}^d$ be the vector function

$$\Lambda_\mu f(x) = \frac{1}{\omega_d} \int_{\mathbb{R}^d} \frac{(y-x) f^\mu(y)}{\|x-y\|^d} \mu(dy),$$

where ω_d is the surface area of the d -dimensional sphere S_{d-1} . If $d = 1$, then $\Lambda_\mu f$ has several equivalent representations

$$\begin{aligned} \Lambda_\mu f(x) &= \mu(f 1_{[x,\infty)}) - \mu(f) \mu(1_{[x,\infty)}) = \mu(f^\mu 1_{[x,\infty)}) \\ &= \mu(1_{(-\infty,x)}) \mu(f) - \mu(f 1_{(-\infty,x)}) = -\mu(f^\mu 1_{(-\infty,x)}) \end{aligned} \quad (19)$$

When $d = 1$ the function $\Lambda_\mu f$ is well defined for arbitrary $f \in \mathcal{B}(\mathbb{R}^d)$ and $\mu \in \mathcal{P}(\mathbb{R}^d)$. This is not the case when $d > 2$. Let $\mu \in \mathcal{P}(\mathbb{R}^d)$, $d \geq 2$ be a probability measure absolutely continuous with respect to the Lebesgue measure. We say that μ is a p -good probability measure if its density $x \rightarrow d_\mu(x)$ with respect to the Lebesgue measure is locally bounded and p -integrable, where $1 < p < d$. The following proposition gives a necessary condition which insures that $\Lambda_\mu f$ is well defined:

Proposition 1 *If $\mu \in \mathcal{P}(\mathbb{R}^d)$, $d \geq 2$ is a p -good probability measure, then $\Lambda_\mu f$ is well defined.*

Proof. We need to prove that the function

$$y \longrightarrow \frac{y - x}{\|x - y\|^d} (f(y) - \mu(f))$$

is integrable with respect to μ for all $x > 0$. Let $\|f\|$ be the L^∞ -norm of f and $\|\mu|_{B(x,1)}\|$ be the L^∞ -norm of the function $x \rightarrow d_\mu(x)$ restricted to $B(x,1)$, the ball of radius 1 and centre x . Then

$$\begin{aligned} \frac{1}{\omega_d} \int_{\mathbb{R}^d} \left| \frac{y_i - x_i}{\|x - y\|^d} (f(y) - \mu(f)) \right| \mu(dy) &\leq \frac{2\|f\| \|\mu|_{B(x,1)}\|}{\omega_d} \int_{B(x,1)} \frac{1}{\|x - y\|^{d-1}} dy \\ &+ \int_{B(x,1)^c} \frac{1}{\|x - y\|^{d-1}} \mu(dy), \end{aligned} \quad (20)$$

where $B(x,1)^c \triangleq \mathbb{R}^d \setminus B(x,1)$. By using polar coordinates it is easy to check that the first integral is equal to ω_d and hence it is finite. Using Hölder's inequality and polar coordinates, we get that

$$\begin{aligned} \int_{B(x,1)^c} \frac{1}{\|x - y\|^{d-1}} \mu(dy) &\leq \left(\int_{B(x,1)^c} \frac{1}{\|x - y\|^{q(d-1)}} dy \right)^{\frac{1}{q}} \left(\int_{B(x,1)^c} (d_\mu(y))^p dy \right)^{\frac{1}{p}} \\ &\leq \left(\omega_d \int_1^\infty r^{-(q-1)(d-1)} dr \right)^{\frac{1}{q}} \|\mu\|_p \end{aligned}$$

and since $(q-1)(d-1) = \frac{d-1}{p-1} > 1$ we get that $\int_1^\infty r^{-(q-1)(d-1)} dr$ is finite. Hence the second integral on the right hand side of (20) is finite and so $\Lambda_\mu f$ is well defined and is a bounded function. ■

We now define the first order differential operators

$$L_\mu^f \varphi(x) \triangleq \sum_{j=1}^d \Lambda_\mu^j f(x) \partial_{x_j} \varphi(x), \quad \bar{L}_\mu^f \varphi(x) \triangleq \frac{L_\mu^f \varphi(x)}{d_\mu(x)}.$$

In the following, we will denote by l_d to be the Lebesgue measure on $\mathbb{R}^d$.

Proposition 2

i. If $\varphi \in \mathcal{C}_b^1(\mathbb{R})$, then

$$\mu(\varphi f^\mu) = l_1(L_\mu^f \varphi) = \mu(\bar{L}_\mu^f \varphi).$$

ii. If $d \geq 2$, $\varphi \in \mathcal{C}_k^1(\mathbb{R}^d)$ and $\mu \in \mathcal{P}(\mathbb{R}^d)$, $d \geq 2$ is a p -good probability measure ($1 < p < d$), then

$$\mu(\varphi f^\mu) = l_d(L_\mu^f \varphi) = \mu(\bar{L}_\mu^f \varphi).$$

Proof.

i. Using the representation (19) we have that $\lim_{|x| \rightarrow \infty} \Lambda_\mu^j f(x) = 0$, hence by integration by parts

$$\int_{\mathbb{R}} \varphi(x) f^\mu(x) \mu(dx) = \int_{\mathbb{R}} \varphi' \Lambda_\mu f(x) dx = \int_{\mathbb{R}} L_\mu^f \varphi(x) dx = \int_{\mathbb{R}} \bar{L}_\mu^f \varphi(x) \mu(dx).$$

ii. For $\varphi \in \mathcal{C}_1^k(\mathbb{R}^d)$ the following identity holds true (see, for example, [26] page 12)

$$\varphi(y) = \frac{1}{\omega_d} \int_{\mathbb{R}^d} \frac{\langle y - x, \nabla \varphi(x) \rangle}{\|x - y\|^d} dx.$$

We will show that the function

$$(x, y) \in \mathbb{R}^d \times \mathbb{R}^d \longrightarrow \frac{\langle y - x, \nabla \varphi(x) \rangle}{\|x - y\|^d} f^\mu(y) d_\mu(y)$$

is integrable with respect to l_{2d} . Hence by Fubini's theorem

$$\begin{aligned} \mu(\varphi f^\mu) &= \int_{\mathbb{R}^d} \varphi(y) f^\mu(y) d_\mu(y) dy \\ &= \int_{\mathbb{R}^d} \frac{1}{\omega_d} \int_{\mathbb{R}^d} \frac{\langle y - x, \nabla \varphi(x) \rangle}{\|x - y\|^d} dx f^\mu(y) d_\mu(y) dy \\ &= \int_{\mathbb{R}^d} \left\langle \frac{1}{\omega_d} \int_{\mathbb{R}^d} \frac{(y - x) f^\mu(y)}{\|x - y\|^d} d_\mu(y) dy, \nabla \varphi(x) \right\rangle \\ &= \int_{\mathbb{R}^d} \langle \Lambda_\mu f(x), \nabla \varphi(x) \rangle = \int_{\mathbb{R}^d} L_\mu^f \varphi(x) dx. \end{aligned}$$

First let us observe that

$$\left| \frac{\langle y - x, \nabla \varphi(x) \rangle}{\|x - y\|^d} f^\mu(y) d_\mu(y) \right| \leq 2 \|f\| \frac{\|\nabla \varphi(x)\|}{\|x - y\|^{d-1}} d_\mu(y).$$

Let now I_1 be the following Riesz type operation

$$I_1\psi(y) = \int_{\mathbb{R}^d} \frac{\psi(x)}{\|x-y\|^{d-1}} dx$$

for $\psi \in \mathcal{C}^k(\mathbb{R}^d)$. Theorem 1.2.1 from [26], page 12, states that $I_1\psi$ is q -integrable for any q such that

$$\frac{1}{q} = \frac{1}{p'} - \frac{1}{d} \text{ where } 1 < p' < d.$$

So $I_1\psi$ is q -integrable for any q such that $0 < \frac{1}{q} < 1 - \frac{1}{d}$. Hence $I_1\psi$ is q -integrable for q such that $\frac{1}{q} = 1 - \frac{1}{p}$. So, by Fubini's theorem (for non-negative functions) and Hölder's inequality,

$$\begin{aligned} \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{\|\nabla\varphi(x)\|}{\|x-y\|^{d-1}} d_\mu(y) dx dy &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \frac{\|\nabla\varphi(x)\|}{\|x-y\|^{d-1}} dx \right) d_\mu(y) dy \\ &= \int_{\mathbb{R}^d} I_1\psi(y) d_\mu(y) dy \\ &\leq \left(\int_{\mathbb{R}^d} I_1\psi(y)^q dy \right)^{\frac{1}{q}} \left(\int_{\mathbb{R}^d} d_\mu(y)^p dy \right)^{\frac{1}{p}} < \infty, \end{aligned}$$

where $\psi(x) \equiv \|\nabla\varphi(x)\|$. Hence our claim. ■

We are now ready to define the coefficients $\tilde{b}_s$ of the equation (9). Let μ be a probability measure that satisfies the following conditions:

1. μ is absolutely continuous with respect to the Lebesgue measure and its density $x \rightarrow d_\mu(x)$ with respect to the Lebesgue measure is a strictly positive function in $C_b^1(\mathbb{R}^d)$.

2. If $d \geq 2$, then there exists p such that μ is a p -good probability measure.

Then the following coefficients are well defined

$$(s, \mu, x) \longrightarrow \tilde{b}_s^\delta(\mu, x) \triangleq b(x) + \frac{\Lambda_\mu \alpha_s^\delta(x)}{d_\mu(x)}, j = 1, \dots, m. \quad (21)$$

Let $X = (X_t)_{t \geq 0}$ be a continuous d -dimensional stochastic process. In the following we will denote by $\tilde{\vartheta}_t$ the distribution of X_t . We say that X is a *good* process if the following three conditions are satisfied:

- $\tilde{\vartheta}_t$ is absolutely continuous with respect to the Lebesgue measure and its density $x \rightarrow \tilde{\vartheta}_t(x)$ is positive for all $x \in \mathbb{R}^d$.

- The function $(t, x) \in [0, \infty] \times \mathbb{R}^d \longrightarrow \tilde{\vartheta}_t(x)$ is continuous.
- If $d \geq 2$, then there exists p such that $\tilde{\vartheta}_t$ is a p -good probability measure for all $t \geq 0$.

We have the following proposition

Proposition 3 *Let $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ be a probability space on which there exists a good process $X = (X_t^\delta)_{t \geq 0}$ which satisfies equation (9). That is*

$$X_t^\delta = X_0^\delta + \int_0^t \tilde{b}_s^\delta \left(\tilde{\vartheta}_s^\delta, X_s^\delta \right) ds + \int_0^t \sigma_s \left(X_s^\delta \right) dB_s$$

where the coefficients $\tilde{b}_s$ are those specified in (21), $B = (B_t)_{t \geq 0}$ be a d -dimensional Brownian motion and X_0^δ has distribution $\bar{\vartheta}_0$. Then $\tilde{\vartheta}_t^\delta$ will be equal to $\bar{\vartheta}_t^\delta$.

Proof. Let us apply Itô's formula to the equation (9) for $\varphi \in C_k^2(\mathbb{R}^d)$. We get that

$$\varphi(X_t^\delta) = \varphi(X_0^\delta) + \int_0^t \left(L_s \varphi(X_s^\delta) + \bar{L}_{\tilde{\vartheta}_s^\delta}^{\alpha_s^\delta} \varphi(X_s^\delta) \right) ds + \int_0^t \sum_{i,j=1}^d \sigma_s^{ij}(X_s) \partial_{x_i} \varphi(X_s) dB_s^j,$$

which yields, by taking expectation and applying Proposition 2, that $\tilde{\vartheta}^\delta$ satisfies

$$\begin{aligned} \tilde{\vartheta}_t^\delta(\varphi) &= \tilde{\vartheta}_0^\delta(\varphi) + \int_0^t \tilde{\vartheta}_s^\delta \left(\left(L_s + \bar{L}_{\tilde{\vartheta}_s^\delta}^{\alpha_s^\delta} \right) \varphi \right) ds \\ &= \tilde{\vartheta}_0^\delta(\varphi) + \int_0^t \tilde{\vartheta}_s^\delta \left(\alpha_s^\delta \varphi + L_s \varphi \right) - \tilde{\vartheta}_s^\delta \left(\alpha_s^\delta \right) \tilde{\vartheta}_s^\delta(\varphi) ds \end{aligned}$$

So $\tilde{\vartheta}^\delta$ satisfied the PDE (17) and, using the uniqueness of the solution of (17), it follows that $\tilde{\vartheta}^\delta = \bar{\vartheta}^\delta$. ■

4 Uniqueness of the Solution

In the following we will prove the uniqueness of a solution of (9) in the class of good of processes as defined in the previous section.

Theorem 4 *There exists at most one solution of (9) which is a good process.*

Proof. First, we note that if $X^\delta = (X_t^\delta)_{t \geq 0}$ is a solution of (9) then the distribution of X_t^δ satisfies (17) and therefore is uniquely determined and equal to $\bar{v}_t^\delta$. Therefore we only need to justify the uniqueness of the solution of

$$dX_t^{i,\delta} = \tilde{b}_t^\delta \left(\bar{v}_t^\delta, X_t^{i,\delta} \right) dt + \sigma_t \left(X_t^i \right) dB_t^i, \quad (22)$$

which is obtained from (9) by replacing $\tilde{v}_t^\delta$ with $\bar{v}_t^\delta$. Hence (22) is an ordinary SDE whose coefficients are locally Lipschitz. In particular $x \rightarrow \frac{1}{\vartheta_t(x)}$ is locally Lipschitz. Hence, for example by Theorem 3.1 page 164 in [13], there exists a stopping time ζ such that equation (9) has a unique solution in the interval $[0, \zeta)$ and, on the event, $\zeta < \infty$ we have

$$\limsup_{t \rightarrow \zeta} \|X_t^\delta\| = \infty.$$

We want to prove next that the event $\zeta < \infty$ has null probability (hence the solution is unique for all $t > 0$). To do this, it is enough to show that, for all $t > 0$ we have $\|X_t^\delta\| < \infty$ P -almost surely. This fact implies, by the continuity of the trajectories, that $\sup_{[0,t]} \|X_s^\delta\| < \infty$ almost surely and hence that $\zeta \geq t$ almost surely. For this it suffices to prove that

$$\lim_{k \rightarrow \infty} P \left(\|X_t^\delta\| \geq k \right) = 0. \quad (23)$$

We have

$$P \left(\|X_t^\delta\| \geq k \right) = \tilde{v}_t^\delta(\varphi_k) = \bar{v}_t^\delta(\varphi_k) = \frac{\vartheta_t^\delta(\varphi_k)}{\vartheta_t^\delta(1)} \leq \frac{\vartheta_t^\delta(\bar{\varphi}_k)}{\vartheta_t^\delta(1)},$$

where

$$\varphi_k(x) \triangleq I_{B(0,k)}(x) = \begin{cases} 1 & \text{if } \|x\| \geq k \\ 0 & \text{otherwise.} \end{cases}, \quad \bar{\varphi}_k(x) \triangleq \begin{cases} 1 & \text{if } \|x\| \geq k \\ \exp\left(\frac{\|x\| - k}{\|x\| - \frac{k}{2}}\right) & \text{if } \frac{k}{2} < \|x\| < k \\ 0 & \text{if } \|x\| < \frac{k}{2} \end{cases}$$

So (23) is implied by

$$\lim_{k \rightarrow \infty} \vartheta_t^\delta(\bar{\varphi}_k) = 0. \quad (24)$$

Now since the coefficients of the operator L_s are all uniformly bounded it follows that $\bar{\varphi}_k \in \bigcap_{s \geq 0} \mathcal{D}(L_s)$ and that $\lim_{k \rightarrow \infty} \sup_{s \in [0, \infty)} \|L_s \bar{\varphi}_k\| = 0$. For arbitrary $T > 0$, define now the function $\Psi : [0, T] \rightarrow \mathbb{R}_+$

$$\Psi(t) = \sup_{s \in [0, t]} \vartheta_t^\delta(\bar{\varphi}_k).$$

From (17) one deduces that, for arbitrary $T > 0$ and $t \in [0, T]$ we have

$$\Psi(t) \leq \bar{\vartheta}_0(\bar{\varphi}_k) + \sup_{s \in [0, T]} \|L_s \bar{\varphi}_k\| \int_0^T \vartheta_s^\delta(1) ds + K_{T, \alpha, \beta, \omega}^\delta \int_0^t \Psi(s) ds,$$

where

$$\begin{aligned} K_{T, \alpha, \beta, \omega}^\delta &= \sup_{s \in [0, T]} \|\alpha_s\| \\ &+ \frac{1}{2} \sum_{j=1}^m \left(\sup_{s \in [0, T]} \|\beta_s^j\|^2 + \frac{1}{\delta} \sup_{s \in [0, T]} \|\beta_s^j\| \max_{i=0, 1, \dots, \lceil \frac{T}{\delta} \rceil + 1} (W_{t_{i+1}}^j(\omega) - W_{t_i}^j(\omega)) \right) \end{aligned}$$

Hence, by Gronwall's inequality,

$$\sup_{s \in [0, t]} \vartheta_s^\delta(\bar{\varphi}_k) \leq e^{K_{T, \alpha, \beta, \omega}^\delta T} \left(\bar{\vartheta}_0(\bar{\varphi}_k) + \sup_{s \in [0, T]} \|L_s \bar{\varphi}_k\| \int_0^T \vartheta_s^\delta(1) ds \right)$$

which implies that

$$\lim_{k \rightarrow \infty} \sup_{s \in [0, t]} \vartheta_s^\delta(\bar{\varphi}_k) = 0. \quad (25)$$

as $\lim_{k \rightarrow \infty} \bar{\vartheta}_0(\bar{\varphi}_k) = 0$. But (25) implies (24), which in turn implies (23) and that completes the proof of the Theorem. ■

5 Existence of the Solution

We are now ready to complete the last step of the programme. For this we need to add one final condition:

ES: Let $(s, x) \longrightarrow \tilde{b}_s^{\delta, \bar{\vartheta}_s^\delta}(x)$ be the function defined in (21) where the measure μ is chosen to be $\bar{\vartheta}_s^\delta$. In other words,

$$\tilde{b}_s^{\delta, \bar{\vartheta}_s^\delta}(x) \triangleq b_s(x) + \frac{\Lambda_{\bar{\vartheta}_s^\delta} \alpha_s^\delta(x)}{\bar{\vartheta}_s^\delta(x)} \quad (26)$$

We will assume that $(s, x) \longrightarrow \tilde{b}_s^{\delta, \bar{\vartheta}_s^\delta}(x)$ is globally Lipschitz. More precisely we will assume that, for any $T \geq 0$, there exists a constant K_T such that

$$\left| \tilde{b}_s^{\delta, \bar{\vartheta}_s^\delta}(x) - \tilde{b}_s^{\delta, \bar{\vartheta}_s^\delta}(y) \right| \leq K_T \|x - y\| \quad (27)$$

for all $x, y \in R^d$ and $s \in [0, T]$.

Theorem 5 *Under the conditions **EU+PD+ES**, equation (9) has a solution which is a good process.*

Proof. We prove the existence of the solution on an arbitrary time interval $[0, T]$. From (27) and the fact that the function

$$s \in [0, T] \longrightarrow b_s(0) + \frac{\Lambda_{\bar{\vartheta}_s^\delta} \alpha_s^\delta(0)}{\bar{\vartheta}_s^\delta(0)}$$

is continuous, hence bounded, it follows (for example, by using Theorem 2.9, p. 289, in [14]) that the equation

$$d\tilde{X}_t^{i,\delta} = \tilde{b}_s^{\delta, \bar{\vartheta}_s^\delta}(\tilde{X}_t^{i,\delta}) dt + \sigma_t(\tilde{X}_t^i) dB_t^i, \quad (28)$$

has a (unique) solution whose distribution satisfies the linear PDE

$$\tilde{\vartheta}_t^\delta(\varphi) = \bar{\vartheta}_0(\varphi) + \int_0^t \tilde{\vartheta}_s^\delta \left(L_s \varphi + \bar{L}_{\bar{\vartheta}_s^\delta}^{\alpha_s^\delta} \varphi \right) ds \quad (29)$$

which has a unique solution (see ... in [18]). From (15) it follows that $\bar{\vartheta}_s^\delta$ is a solution of PDE (29), hence $\tilde{\vartheta}^\delta$ and $\bar{\vartheta}^\delta$, hence $\tilde{X}^{i,\delta}$ is in fact a solution of the nonlinear SDE (9). ■

Remark 1 *If $d = 1$ and for all $s \geq 0$, the density of ϑ_s^δ with respect to the Lebesgue measure has the form*

$$\vartheta_s^\delta(x) = e^{-F_s^\delta(x)}, \quad x \in \mathbb{R}.$$

*where F_s^δ is a differentiable convex function, condition **ES** is satisfied.*

Proof. We have three cases:

1. If $\frac{dF_s^\delta}{dx}(x) = 0$, then $\frac{d}{dx} \left(\frac{\Lambda_{\vartheta_s^\delta} \alpha_s^\delta(x)}{e^{-F_s^\delta(x)}} \right) = (\alpha_s^\delta)^\mu(x)$ which is bounded since α_s^δ is bounded.
2. If $\frac{dF_s^\delta}{dx}(x) > 0$, then using the first part of the representation (19) we get

$$\frac{d}{dx} \left(\frac{\Lambda_{\vartheta_s^\delta} f(x)}{e^{-F_s^\delta(x)}} \right) = (\alpha_s^\delta)^\mu(x) + \int_x^\infty (\alpha_s^\delta)^\mu(y) \frac{dF_s^\delta}{dx}(x) e^{F_s^\delta(x) - F_s^\delta(y)} dy$$

Since F_s^δ is convex, for $y > x$ we have

$$\frac{F_s^\delta(y) - F_s^\delta(x)}{y - x} \geq \frac{dF_s^\delta}{dx}(x) \implies F_s^\delta(x) - F_s^\delta(y) \leq (x - y) \frac{dF_s^\delta}{dx}(x)$$

So

$$\begin{aligned} \left| \int_x^\infty (\alpha_s^\delta)^\mu(y) \frac{dF_s^\delta}{dx}(x) e^{F_s^\delta(x) - F_s^\delta(y)} dy \right| &\leq \|(\alpha_s^\delta)^\mu\| \frac{dF_s^\delta}{dx}(x) \int_x^\infty e^{(x-y) \frac{dF_s^\delta}{dx}(x)} dy \\ &\leq \|(\alpha_s^\delta)^\mu\| \end{aligned}$$

since $\frac{dF_s^\delta}{dx}(x) \int_x^\infty e^{(x-y) \frac{dF_s^\delta}{dx}(x)} dy = 1$.

3. If $\frac{dF_s^\delta}{dx}(x) < 0$, then then using the second part of the representation (19) we get

$$\frac{d}{dx} \left(\frac{\Lambda_{\partial_s^\delta} f(x)}{e^{-F_s^\delta(x)}} \right) = (\alpha_s^\delta)^\mu(x) - \int_{-\infty}^x (\alpha_s^\delta)^\mu(y) \frac{dF_s^\delta}{dx}(x) e^{F_s^\delta(x) - F_s^\delta(y)} dy$$

and we follow the same steps as in the previous case. ■

We complete the section by noting that the above construction works with minimal changes when (the initial Markov process) ξ is a reflecting boundary diffusion. In this case, the analysis simplifies considerably if the domain is compact. For example, the cumbersome condition **ES** is replaced by the assumption that ξ has a density which is bounded away from 0. We will detail the analysis of this case in a forthcoming paper, together with the description of the associated numerical algorithm. However, for completeness, we briefly present the results here:

Following the notation and results in [19], assume that ξ is the solution of a *stochastic differential equation with reflection along the normal*. In other words, ξ is the unique solution of the equation

$$\xi_t = \xi_0 + \int_0^t b_s(\xi_s) ds + \int_0^t \sigma_s(\xi_s) dB_s - k_t, \quad (30)$$

where $\xi_t \in \overline{B(0, M)^2}$ for all $t \geq 0$ and k is a bounded variation process

$$|k|_t = \int_0^t 1_{\{\xi_s \in \partial B(0, M)\}} d|k|_s, \quad k_t = \int_0^t \frac{\xi_s}{M} d|k|_s$$

For general domains D , k is defined as

$$|k|_t = \int_0^t 1_{\{\xi_s \in \partial D\}} d|k|_s, \quad k_t = \int_0^t n(\xi_s) d|k|_s$$

where $n(x)$ is the unit outward normal to ∂D at x .

Obviously the generator $L = (L_s)_{s \geq 0}$ associated to ξ has the form (2) for any φ in $C_K^2(B(0, M))$, the set of twice differentiable functions defined on $B(0, M)$

² $B(0, M)$ is the ball of center 0 and radius M .

with compact support. The operator $\Lambda_\mu f$ is well defined for measures μ with support in $\overline{B(0, M)}$ absolute continuous with respect to the Lebesgue measure and with bounded density. The definition of a *good* process is now slightly simpler: A continuous process $X = (X_t)_{t \geq 0}$ which takes values in $\overline{B(0, M)}$ is a *good* process if the following two conditions are satisfied (as before, $\tilde{\vartheta}_t$ is the distribution of X_t):

◆ $\tilde{\vartheta}_t$ is absolutely continuous with respect to the Lebesgue measure and its density $x \rightarrow \tilde{\vartheta}_t(x)$ is positive on $\overline{B(0, M)}$.

◆ The function $(t, x) \in [0, \infty] \times \overline{B(0, M)} \rightarrow \tilde{\vartheta}_t(x)$ is continuous and bounded.

Proposition (3) now becomes:

Proposition 6 *Let $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ be a probability space on which there exists a good process $X^\delta = (X_t^\delta)_{t \geq 0}$ which satisfies equation*

$$X_t^\delta = X_0^\delta + \int_0^t \tilde{b}_s^\delta(\tilde{\vartheta}_s^\delta, X_s^\delta) ds + \int_0^t \sigma_s(X_s^\delta) dB_s - k_t^\delta \quad (31)$$

where $X_t^\delta \in \overline{B(0, M)}$ for all $t \geq 0$ and k^δ is a bounded variation process

$$|k^\delta|_t = \int_0^t 1_{\{X_s^\delta \in \partial B(0, M)\}} d|k^\delta|_s, \quad k_t^\delta = \int_0^t \frac{X_s^\delta}{M} d|k^\delta|_s$$

In (31), the coefficients $\tilde{b}_s$ are those specified in (21), $B = (B_t)_{t \geq 0}$ be a d -dimensional Brownian motion and X_0^δ has distribution $\tilde{\vartheta}_0$. Then the distribution of X_t^δ will be equal to $\tilde{\vartheta}_t^\delta$.

The uniqueness of the pair (X^δ, k^δ) follows by slight variation of the argument for the proof of theorem (4). The existence of the pair (X^δ, k^δ) requires the following simpler condition:

ES': Assume that, $\xi = (\xi_t)_{t \geq 0}$, the solution of the SDE (30) is a good process and that its density $\check{\vartheta}_t(x)$ is bounded away from 0. More precisely, for any $T > 0$ we have

$$\inf_{t \in [0, T] x \in \overline{B(0, M)}} \check{\vartheta}_t(x) > 0.$$

Then we have the following result:

Theorem 7 *Under the conditions **EU+PD+ES'**, equation (31) has a unique solution which is a good process.*

The results follows by proving that **ES'** implies **ES**. Since $\bar{\vartheta}_t^\delta$ has the representation

$$\bar{\vartheta}_t^\delta(\varphi) = E \left[\varphi(\xi_t) \exp \left(\int_0^t \alpha_s^\delta(\xi_s) - \bar{\vartheta}_t^\delta(\alpha_s^\delta) ds \right) \right]. \quad (32)$$

$\bar{\vartheta}_t^\delta$ is absolutely continuous with respect to the Lebesgue measure, too. Further, since α_s^δ as defined in (16) is a bounded function on $\overline{B(0, M)}$, we deduce that there exists a positive constant ϵ such that

$$\bar{\vartheta}_t^\delta(\varphi) \geq \epsilon E[\varphi(\xi_t)] = \epsilon \int_{B(0, M)} \varphi(x) \check{\vartheta}_t(x) dx$$

Hence the density of $\bar{\vartheta}_t^\delta$ is uniformly bounded away from 0. One can also prove that the density of $\bar{\vartheta}_t^\delta$ has uniformly bounded first order derivatives. This implies that the function $(s, x) \longrightarrow \frac{\Lambda_{\bar{\vartheta}_s^\delta} \alpha_s^\delta(x)}{\check{\vartheta}_s^\delta(x)}$ is globally Lipschitz, hence **ES** is satisfied.

Remark 2 *Numerical methods for equations of type (30) have extensively developed (see for example [23], [24] and the references therein). We will adapt these schemes in order to produce a numerical method for solving (31).*

6 Rates of convergence and final remarks

It is obvious that, for any $\varphi \in C_b(\mathbb{R}^d)$, we have

$$\tilde{E} \left[\left(\bar{\vartheta}_t^\delta(\varphi) - \bar{\vartheta}_t^{\delta, n}(\varphi) \right)^4 \right] \leq \frac{\|\varphi\|^4}{n^2}$$

so

$$\tilde{E} \left[\left\| \bar{\vartheta}_t^\delta - \bar{\vartheta}_t^{\delta, n} \right\|_{\mathcal{M}}^4 \right] \leq \frac{1}{n^2}.$$

where the norm $\|\cdot\|_{\mathcal{M}}$ is defined in (14). From (13), we have that

$$\left\| \vartheta_t^{\delta(\omega, n)} - \vartheta_t \right\|_{\mathcal{M}} \leq \frac{K}{\sqrt{n}} \text{ and } \left\| \bar{\vartheta}_t^{\delta(\omega, n)} - \bar{\vartheta}_t \right\|_{\mathcal{M}} \leq \frac{\bar{K}}{\sqrt{n}},$$

hence

$$\tilde{E} \left[\left\| \bar{\vartheta}_t - \bar{\vartheta}_t^{\delta(\omega, n), n} \right\|_{\mathcal{M}}^4 \right] \leq \frac{\left(\sqrt[4]{\bar{K}} + 1 \right)^4}{n^2},$$

so $\bar{\vartheta}_t^{\delta(\omega,n),n}$ converges to $\bar{\vartheta}_t$ $\tilde{P}$ -almost surely. We also have almost sure convergence if we view $\bar{\vartheta}_t^{\delta(\omega,n),n}$ and $\bar{\vartheta}_t$ as processes on the product space $(\bar{\Omega}, \bar{\mathcal{F}}, \bar{P})$

$$(\bar{\Omega}, \bar{\mathcal{F}}, \bar{P}) = (\Omega \times \hat{\Omega}, \mathcal{F} \otimes \mathcal{F}, \tilde{P} \otimes \tilde{P})$$

on which we ‘lift’ the processes $\bar{\vartheta}_t^{\delta(\omega,n),n}$ and $\bar{\vartheta}_t$ from the component spaces. Finally, in introduction we chose the asymptotic representation for ϑ_t to be

$$\vartheta_t = \lim_{n \rightarrow \infty} \frac{a_t^n}{n} \sum_{i=1}^n \delta_{X_t^{i,\delta(W,(\omega),n)}}, \quad (33)$$

where

$$\begin{aligned} a_t^n = & \exp \left(\int_0^t \bar{\vartheta}_t^{\delta(\omega,n),n} (\alpha_s) ds + \sum_{j=1}^m \int_0^t \bar{\vartheta}_t^{\delta(\omega,n),n} (\beta_s^j) d(W_s^{\delta(\omega,n)})^j \right. \\ & \left. - \frac{1}{2} \sum_{j=1}^m \int_0^t \bar{\vartheta}_t^{\delta(\omega,n),n} (\beta_s^j)^2 ds \right) \end{aligned} \quad (34)$$

Then each of the terms in the formula (34), i.e., $\int_0^t \bar{\vartheta}_t^{\delta(\omega,n),n} (\alpha_s) ds$, $\int_0^t \bar{\vartheta}_t^{\delta(\omega,n),n} (\beta_s^j) d(W_s^{\delta(\omega,n)})^j$ and $\int_0^t \bar{\vartheta}_t^{\delta(\omega,n),n} (\beta_s^j)^2 ds$ converge $\bar{P}$ -almost surely to $\int_0^t \bar{\vartheta}_t (\alpha_s) ds$, $\int_0^t \bar{\vartheta}_t (\beta_s^j) dW_s^j$ and, respectively, $\int_0^t \bar{\vartheta}_t (\beta_s^j)^2 ds$, hence a_t^n converges almost surely to $\vartheta_t(1)$, so (34) holds true.

References

- [1] J. M. C. CLARK, *The design of robust approximations to the stochastic differential equations of nonlinear filtering*. Communication systems and random process theory (Proc. 2nd NATO Advanced Study Inst., Darlington, 1977), pp. 721–734. NATO Advanced Study Inst. Ser., Ser. E: Appl. Sci., No. 25, Sijthoff & Noordhoff, Alphen aan den Rijn, 1978.
- [2] J. M. C. CLARK, D. CRISAN, *On a robust version of the integral representation formula of nonlinear filtering*, to appear in Probability Theory and Related Fields, 2005.
- [3] D. CRISAN, *Numerical methods for solving the stochastic filtering problem*, Numerical methods and stochastics (Toronto, ON, 1999), 1–20, Fields Inst. Commun., 34, Amer. Math. Soc., Providence, RI, 2002.

- [4] D. CRISAN, J. GAINES, T. LYONS, *Convergence of a branching particle method to the solution of the Zakai equation*. SIAM J. Appl. Math. 58 (1998), no. 5, 1568–1590 (electronic).
- [5] D. CRISAN, T. LYONS, *A particle approximation of the solution of the Kushner-Stratonovitch equation* Probab. Theory Related Fields 115 (1999), no. 4, 549–578.
- [6] D. CRISAN, P. DEL MORAL T. LYONS, *Interacting particle systems approximations of the Kushner-Stratonovitch equation*. Adv. in Appl. Probab. 31 (1999), no. 3, 819–838.
- [7] D. CRISAN, *Particle filters—a theoretical perspective*. Sequential Monte Carlo methods in practice, 17–41, Stat. Eng. Inf. Sci., Springer, New York, 2001.
- [8] D. CRISAN, A. DOUCET, *A survey of convergence results on particle filtering methods for practitioners*. IEEE Trans. Signal Process. 50, no. 3, pp 736–746, 2002.
- [9] D. CRISAN, *Exact rates of convergence for a branching particle approximation to the solution of the Zakai equation* Ann. Probab. 31 (2003), no. 2, 693–718.
- [10] P. DEL MORAL, L. MICLO, *Branching and interacting particle systems approximations of Feynman-Kac formulae with applications to non-linear filtering*. Séminaire de Probabilités, XXXIV, 1–145, Lecture Notes in Math., 1729, Springer, Berlin, 2000.
- [11] T. E. DUNCAN, *Likelihood functions for stochastic signals in white noise*, Information and Control 16 1970 303–310.
- [12] M. FUJISAKI, G. KALLIANPUR, H. KUNITA, *Stochastic differential equations for the non linear filtering problem*. Osaka J. Math. 9, pp 19–40, 1972.
- [13] N. IKEDA, S. WATANABE, *Stochastic differential equations and diffusion processes*, North-Holland Mathematical Library, 24. North-Holland Publishing Co., Amsterdam-New York; Kodansha, Ltd., Tokyo, 1981.
- [14] I. KARATZAS, S. E. SHREVE, *Brownian motion and stochastic calculus*, Second Edition. Graduate Texts in Mathematics, 113. Springer-Verlag, New York, 1991.
- [15] T. G. KURTZ, J. XIONG, *Particle Representation for a Class of Nonlinear SPDE's*, Stochastic Process. Appl. 83, no. 1, pp 103–126, 1999.

- [16] T. G. KURTZ, J. XIONG, *Numerical Solutions for a Class of SPDEs with Application to Filtering*. Stochastics in finite and infinite dimensions, 233–258, Trends Math., Birkhäuser Boston, 2001.
- [17] H. J. KUSHNER, *Dynamical equations for optimal nonlinear filtering*, J. Differential Equations 3, 179–190, 1967.
- [18] O. A. LADYŽENSKAJA, V. A. SOLONNIKOV, N. N. URAL’CEVA, *Linear and quasilinear equations of parabolic type*. (Russian) Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23 American Mathematical Society, Providence, R.I. 1967.
- [19] P. L. LIONS, A.S. SZNITMAN, *Stochastic differential equations with reflecting boundary conditions*, Comm. Pure Appl. Math. 37, no. 4, 511–537, 1984.
- [20] R. S. LIPTSER, A. N. SHIRYAEV, *Statistics of random processes. I. General theory*, Second, revised and expanded edition, Springer-Verlag, Berlin, 2001.
- [21] R. E. MORTENSEN, *Stochastic optimal control with noisy observations*, Internat. J. Control (1) 4 1966 455–464.
- [22] E. PARDOUX, *Filtrage Non Linéaire et Equations aux Dérivées Partielles Stochastiques Associées*, Ecole d’Eté de Probabilités de Saint-Flour XIX - 1989, Lecture Notes in Mathematics, 1464, Springer-Verlag, 1991.
- [23] R. PETTERSSON, *Penalization schemes for reflecting stochastic differential equations*, Bernoulli 3, no. 4, 403–414, 1997.
- [24] L. SLOMINSKI, *Euler’s approximations of solutions of SDEs with reflecting boundary*, Stochastic Process. Appl. 94, no. 2, 317–337, 2001.
- [25] B. L. ROZOVSKII, *Stochastic Evolution Systems*, Kluwer, Dordrecht, 1990.
- [26] L. SALOFF-COSTE, *Aspects of Sobolev-type inequalities*, London Mathematical Society Lecture Note Series, 289. Cambridge University Press, Cambridge, 2002.
- [27] M ZAKAI, *On the optimal filtering of diffusion processes*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, 11, pp 230–243, 1969.